

BOTANY :: 1999

1. The earlier institute established at Pusa, is now called as
1) BIs 2) ICAR 3) IARi 4) CsIR
2. Balancing roots are seen in
(1) Dolichos (2) Pistia (3) Piper (4) Vanda
3. Root, stem and leaf are all modified in
(1) Dracaena (2) Aloe (3) Asparagus (4) Opuntia
4. The following features is found in both Balanophora and Casuarina
(1) Parasitism (2) Tree habit (3) Scale leaves (4) Phyllode
5. The following plant produces both offsets and bulbils
(1) Chrysanthemum (2) Oxalis (3) Agave (4) Bryophyllum
6. Inflorescence in Jatropha is a
(1) monochasial scorpioid cyme (2) polychasial cyme
(3) dichasial cyme (4) solitary flower
7. In Fabaceae, the floral structures which enclose the essential organs are
(1) posterior petals (2) lateral petals (3) sepals (4) anterior petals
8. Geitonogamy is found in
(1) Sorghum (2) Erythrina (3) Hydrilla (4) Maize
9. Modification of bract in Typhonium is a
(1) Scale (2) Spathe (3) Involucre (4) Glume
10. Free central placentation is found in
(1) Raphanus (2) Dianthus (3) Dolichos (4) Helianthus
11. The position of the micropyle' found in an orthotropous ovule are turned down about 160° (degrees) with reference to the funicle. In addition the ovule and embryo sac are curved. Such a type of ovule is called as
(1) Anatropous (2) Campylotropous (3) Hemitropous (4) Amphitropous
12. Scale bark is seen in
(1) Terminalia (2) Mangifera (3) Psidium (4) Neem
13. The scientific term which is not related to the primary constriction of a chromosome is
(1) Centromere (2) Kinomere (3) Kinetochore (4) Chromomere
14. Asafoetida is a
(1) Tannin (2) oleoresin (3) latex (4) gum resin
15. The haploid Chromosome number in Saccharum officinale is as many times greater in the following as that of Zeamays
(1) four-fold (2) two-fold (3) six-fold (4) three-fold
16. Tyloses are observed in the lumen of
(1) Tracheids (2) Sieve tubes (3) Fibre tracheids (4) Xylem vessels
17. The genetical importance of raising haploids in higher plants through tissue culture lies their use in the production of
(1) homozygous diploids (2) heterozygous diploids
(3) plants that do not produce flowers (4) hybrid plants
18. As compared to sclerenchyma, parenchyma does NOT function in
(1) gas exchange (2) nutrient transport

36. The Scientist who first carried out experiments' on hybrid vigour in maize was
 (1) Mendle (2) Shull (3) Johannsen (4) Kolreuter
37. An example of planktonic flora of pond ecosystem
 (1) Diatoms (2) Crustaceans (3) Hydrilla (4) Chara
38. Lotic ecosystem refers to
 (1) Static water ecosystem (2) Ecosystem of flowing water
 (3) Ecosystem of estuary water (4) Deep marine water ecosystem
39. The following is a Xerophyte containing air spaces in the stem and roots
 (1) Eicchornia (2) Opuntia (3) Aloe (4) Rhiiphora
40. The following is a primary consumer
 (1) Diatoms (2) Pediastrum (3) Cornivores (4) Rotifers
41. The morphogenetic property of cytokinin was experimentally proved first by ,
 (1) Hanning (2) Guha and Maheswari
 (3) Skoog and Miller (4) Went
42. In a plant physiology experiment a green plant kept in light was found releasing more units of $^{18}\text{O}_2$. Which of the following compounds would have been supplied to that plant?
 (1) $\text{C}_6\text{H}_{12}\text{O}_6$ containing ^{18}O (2) CO_2 containing ^{18}O
 (3) Ozone containing ^{18}O (4) H_2O containing ^{18}O
43. The number of oxidations involving NADH in Krebs cycle is
 (1) 4 (2) 3 (3) 2 (4) 5
44. In noncyclic photophosphorylation, the site of ATP synthesis taking place between the following electron carriers, is
 (1) PS II and Q (2) Cyt. b_6 and Cyt. f
 (3) Cyt. b_6 and plastocyanin (4) PSI and ferredoxin
45. Stomatal closure is induced by
 (1) Indole acetic acid (2) Naphthalene acetic acid
 (3) Indole butyric acid (4) Phenyl mercuric acid
- (1) In its absence, plants fail to reproduce
 (2) It plays a role that is replaceable
 (3) It is necessary at all stages of development
 (4) It affects the metabolism directly
47. Munch Mass flow hypothesis mainly explain the long distance transport of the following
 (1) Mineral salts (2) Sugars (3) Proteins (4) Starch
48. The enzyme code of enzyme 2.7.1.1 refers to the following main group
 (1) Ligase (2) Hydrolase (3) Transferase (4) Lyase
49. The following hormone induces dormancy in potatoes
 (1) IM (2) Gibberellins (3) Absciscic acid (4) Zeatin
50. The following is a constituent of nitrogenase
 (1) magnesium (2) molybdenum. (3) manganese (4) Potassium

ANSWER

(1) 3 (2) 2 (3) 3 (4) 3 (5) 3 |

(6)	3	(7)	4	(8)	4	(9)	2	(10)	2
(11)	4	(12)	3	(13)	4	(14)	4	(15)	1
(16)	4	(17)	1	(18)	3	(19)	3	(20)	3
(21)	3	(22)	4	(23)	3	(24)	1	(25)	2
(26)	2	(27)	2	(28)	3	(29)	4	(30)	3
(31)	2	(32)	1	(33)	3	(34)	1	(35)	1
(36)	2	(37)	1	(38)	2	(39)	4	(40)	4
(41)	3	(42)	4	(43)	2	(44)	2	(45)	Del
(46)	2	(47)	2	(48)	3	(49)	3	(50)	2