

*IIT JEE-2006*

# *Mathematics*

**Time: 2 hours**

**Note:** Question number 1 to 12 carries (3, -1) marks each, 13 to 20 carries (5, -1) marks each, 21 to 32 carries (5, -2) marks each and 33 to 40 carries (6, 0) marks each.

**Section – A (Single Option Correct)**



Sol. (C)

$$\lim_{x \rightarrow 0} \left( (\sin x)^{1/x} - \left(\frac{1}{x}\right)^{\sin x} \right)$$

$$0 + e^{\lim_{x \rightarrow 0} \sin x \ln\left(\frac{1}{x}\right)} = 1 \text{ (using L'Hospital's rule).}$$

2.  $\int \frac{x^2 - 1}{x^3 \sqrt{2x^4 - 2x^2 + 1}} dx$  is equal to

(A)  $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x^2} + c$

(B)  $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x^3} + c$

(C)  $\frac{\sqrt{2x^4 - 2x^2 + 1}}{x} + c$

(D)  $\frac{\sqrt{2x^4 - 2x^2 + 1}}{2x^2} - c$

**Sol.** (D)

$$\int \frac{\left(\frac{1}{x^3} - \frac{1}{x^5}\right) dx}{\sqrt{2 - \frac{2}{x^2} + \frac{1}{x^4}}}$$

$$\text{Let } 2 - \frac{2}{x^2} + \frac{1}{x^4} = z \Rightarrow \frac{1}{4} \int \frac{dz}{\sqrt{z}}$$

$$\Rightarrow \frac{1}{2} \times \sqrt{z} + c \Rightarrow \frac{1}{2} \sqrt{2 - \frac{2}{x^2} + \frac{1}{x^4}} + c$$



**Sol.** (C)

$$\Delta = \frac{\sqrt{3}}{4} b^2 \quad \dots (1)$$

Also  $\frac{\sin 120^\circ}{a} = \frac{\sin 30^\circ}{b} \Rightarrow a = \sqrt{3}b$

and  $\Delta = \sqrt{3}s$  and  $s = \frac{1}{2}(a + 2b)$

$$\Rightarrow \Delta = \frac{\sqrt{3}}{2}(a + 2b) \quad \dots(2)$$

From (1) and (2), we get  $\Delta = (12 + 7\sqrt{3})$ .

4. If  $0 < \theta < 2\pi$ , then the intervals of values of  $\theta$  for which  $2\sin^2\theta - 5\sin\theta + 2 > 0$ , is

- (A)  $\left(0, \frac{\pi}{6}\right) \cup \left(\frac{5\pi}{6}, 2\pi\right)$       (B)  $\left(\frac{\pi}{8}, \frac{5\pi}{6}\right)$   
 (C)  $\left(0, \frac{\pi}{8}\right) \cup \left(\frac{\pi}{6}, \frac{5\pi}{6}\right)$       (D)  $\left(\frac{41\pi}{48}, \pi\right)$

Sol.

(A)

$$\begin{aligned} 2\sin^2\theta - 5\sin\theta + 2 &> 0 \\ \Rightarrow (\sin\theta - 2)(2\sin\theta - 1) &> 0 \\ \Rightarrow \sin\theta &< \frac{1}{2} \\ \Rightarrow \theta &\in \left(0, \frac{\pi}{6}\right) \cup \left(\frac{5\pi}{6}, 2\pi\right). \end{aligned}$$

5. If  $w = \alpha + i\beta$ , where  $\beta \neq 0$  and  $z \neq 1$ , satisfies the condition that  $\left(\frac{w - \bar{w}z}{1 - z}\right)$  is purely real, then the set of values of  $z$  is

- (A)  $\{z : |z| = 1\}$       (B)  $\{z : z = \bar{z}\}$   
 (C)  $\{z : z \neq 1\}$       (D)  $\{z : |z| = 1, z \neq 1\}$

Sol.

(D)

$$\begin{aligned} \frac{w - \bar{w}z}{1 - z} &= \frac{\bar{w} - w\bar{z}}{1 - \bar{z}} \\ \Rightarrow (z\bar{z} - 1)(\bar{w} - w) &= 0 \\ \Rightarrow z\bar{z} = 1 &\Rightarrow |z|^2 = 1 \Rightarrow |z| = 1. \end{aligned}$$

6. Let  $a, b, c$  be the sides of a triangle. No two of them are equal and  $\lambda \in \mathbb{R}$ . If the roots of the equation  $x^2 + 2(a+b+c)x + 3\lambda(ab+bc+ca) = 0$  are real, then

- (A)  $\lambda < \frac{4}{3}$       (B)  $\lambda > \frac{5}{3}$   
 (C)  $\lambda \in \left(\frac{1}{3}, \frac{5}{3}\right)$       (D)  $\lambda \in \left(\frac{4}{3}, \frac{5}{3}\right)$

Sol.

(A)

$$\begin{aligned} D &\geq 0 \\ \Rightarrow 4(a+b+c)^2 - 12\lambda(ab+bc+ca) &\geq 0 \\ \Rightarrow \lambda &\leq \frac{a^2 + b^2 + c^2}{3(ab+bc+ca)} + \frac{2}{3} \\ \text{Since } |a-b| < c &\Rightarrow a^2 + b^2 - 2ab < c^2 \quad \dots(1) \\ |b-c| < a &\Rightarrow b^2 + c^2 - 2bc < a^2 \quad \dots(2) \\ |c-a| < b &\Rightarrow c^2 + a^2 - 2ac < b^2 \quad \dots(3) \end{aligned}$$

From (1), (2) and (3), we get  $\frac{a^2 + b^2 + c^2}{ab+bc+ca} < 2$ .

$$\text{Hence } \lambda < \frac{2}{3} + \frac{2}{3} \Rightarrow \lambda < \frac{4}{3}.$$



$$\therefore \text{Distance of the plane from the point } (1, 2, 2) = \frac{|1+2+1|}{\sqrt{1^2 + 1^2}} = 2\sqrt{2}.$$

12. Let  $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} + \hat{j} - \hat{k}$ . A vector in the plane of  $\vec{a}$  and  $\vec{b}$  whose projection on  $\vec{c}$  is  $\frac{1}{\sqrt{3}}$ , is

- (A)  $4\hat{i} - \hat{j} + 4\hat{k}$       (B)  $3\hat{i} + \hat{j} - 3\hat{k}$   
 (C)  $2\hat{i} - \hat{j} - 2\hat{k}$       (D)  $4\hat{i} + \hat{j} - 4\hat{k}$

**Sol.** (A)

Vector lying in the plane of  $\vec{a}$  and  $\vec{b}$  is  $\vec{r} = \lambda_1 \vec{a} - \lambda_2 \vec{b}$  and its projection on  $\vec{c}$  is  $\frac{1}{\sqrt{3}}$

$$\Rightarrow \left[ (\lambda_1 + \lambda_2) \hat{i} + (2\lambda_1 - \lambda_2) \hat{j} + (\lambda_1 + \lambda_2) \hat{k} \right] \cdot \frac{[\hat{i} + \hat{j} - \hat{k}]}{\sqrt{3}} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow 2\lambda_1 - \lambda_2 = 1 \Rightarrow \bar{r} = (3\lambda_1 - 1)\hat{i} + \hat{j} + (3\lambda_1 - 1)\hat{k}$$

Hence the required vector is  $-2\hat{i} + 5\hat{j} - 2\hat{k}$ .

**Section – B (May have more than one option correct)**



**Sol.** (A), (B)

Equation of tangent to  $x^2 - y$  is

$$y = mx - \frac{1}{4}m^2 \quad \dots(1)$$

Equation of tangent to  $(x - 2)^2 = y$  is

$$y = m(x - 2) + \frac{1}{4}m^2 \quad \dots(2)$$

(1) and (2) are identical.

$$\Rightarrow m = 0 \text{ or } 4$$

$\therefore$  Common tangents are  $y = 0$  and  $y = 4x - 4$ .




**Sol.** (A), (C)

$$f(x) = \min\{1, x^2, x^3\}$$

$$\Rightarrow f(x) = \begin{cases} x^3 & , \quad x \leq 1 \\ 1 & , \quad x > 1 \end{cases}$$

$\Rightarrow f(x)$  is continuous  $\forall x \in R$



15. A tangent drawn to the curve  $y = f(x)$  at  $P(x, y)$  cuts the x-axis and y-axis at A and B respectively such that  $BP : AP = 3 : 1$ , given that  $f(1) = 1$ , then



- (C) curve passes through  $(2, 1/8)$       (D) equation of curve is  $x \frac{dy}{dx} + 3y = 0$

Sol.

(C), (D)

Equation of the tangent is

$$Y - y = \frac{dy}{dx}(X - x)$$


Given  $\frac{BP}{AP} = \frac{3}{1}$  so that

$$\Rightarrow \frac{dx}{x} = -\frac{dy}{3y} \Rightarrow x \frac{dy}{dx} + 3y = 0$$

$$\Rightarrow \ln x = -\frac{1}{3} \ln y - \ln c \Rightarrow \ln x^3 = -(\ln cy)$$

$$\Rightarrow \frac{1}{x^3} = cy. \text{ Given } f(1) = 1 \Rightarrow c = 1$$

$$\therefore y = \frac{1}{x^3}.$$



16.

If a hyperbola passes through the focus of the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  and its transverse and conjugate axes coincide with the major and minor axes of the ellipse, and the product of eccentricities is 1, then

(A) the equation of hyperbola is  $\frac{x^2}{9} - \frac{y^2}{16} = 1$

(B) the equation of hyperbola is  $\frac{x^2}{9} - \frac{y^2}{25} = 1$

(C) focus of hyperbola is  $(5, 0)$

(D) focus of hyperbola is  $(5\sqrt{3}, 0)$

Sol.

(A), (C)

Eccentricity of ellipse =  $\frac{3}{5}$

Eccentricity of hyperbola =  $\frac{5}{3}$  and it passes through  $(\pm 3, 0)$

$\Rightarrow$  its equation  $\frac{x^2}{9} - \frac{y^2}{b^2} = 1$

where  $1 + \frac{b^2}{9} = \frac{25}{9} \Rightarrow b^2 = 16$

$\Rightarrow \frac{x^2}{9} - \frac{y^2}{16} = 1$  and its foci are  $(\pm 5, 0)$ .

17.

Internal bisector of  $\angle A$  of triangle ABC meets side BC at D. A line drawn through D perpendicular to AD intersects the side AC at E and the side AB at F. If a, b, c represent sides of  $\triangle ABC$  then

(A) AE is HM of b and c

(B)  $AD = \frac{2bc}{b+c} \cos \frac{A}{2}$

(C)  $EF = \frac{4bc}{b+c} \sin \frac{A}{2}$

(D) the triangle AEF is isosceles

Sol.

(A), (B), (C), (D).


We have  $\triangle ABC = \triangle ABD + \triangle ACD$ 

$$\Rightarrow \frac{1}{2}bc \sin A = \frac{1}{2}cAD \sin \frac{A}{2} + \frac{1}{2}b \times AD \sin \frac{A}{2}$$

$$\Rightarrow AD = \frac{2bc}{b+c} \cos \frac{A}{2}$$

Again  $AE = AD \sec \frac{A}{2}$

$$= \frac{2bc}{b+c} \Rightarrow AE \text{ is HM of } b \text{ and } c.$$



$$\begin{aligned} EF &= ED + DF = 2 \times AD \tan \frac{A}{2} = \frac{2 \times 2bc}{b+c} \times \cos \frac{A}{2} \times \tan \frac{A}{2} \\ &= \frac{4bc}{b+c} \sin \frac{A}{2} \end{aligned}$$

As  $AD \perp EF$  and  $DE = DF$  and  $AD$  is bisector  $\Rightarrow AEF$  is isosceles.  
Hence A, B, C and D are correct answers.

18.  $f(x)$  is cubic polynomial which has local maximum at  $x = -1$ . If  $f(2) = 18$ ,  $f(1) = -1$  and  $f(x)$  has local minima at  $x = 0$ , then

- (A) the distance between  $(-1, 2)$  and  $(a, f(a))$ , where  $x = a$  is the point of local minima is  $2\sqrt{5}$
- (B)  $f(x)$  is increasing for  $x \in [1, 2\sqrt{5}]$
- (C)  $f(x)$  has local minima at  $x = 1$
- (D) the value of  $f(0) = 5$


**Sol.** (B), (C)

The required polynomial which satisfy the condition

$$\text{is } f(x) = \frac{1}{4} (19x^3 - 57x + 34)$$

$f(x)$  has local maximum at  $x = -1$  and local minimum at  $x = 1$

Hence  $f(x)$  is increasing for  $x \in [1, 2\sqrt{5}]$ .



19. Let  $\vec{A}$  be vector parallel to line of intersection of planes  $P_1$  and  $P_2$  through origin.  $P_1$  is parallel to the vectors  $2\hat{i} + 3\hat{k}$  and  $4\hat{j} - 3\hat{k}$  and  $P_2$  is parallel to  $\hat{j} - \hat{k}$  and  $3\hat{i} + 3\hat{j}$ , then the angle between vectors  $\vec{A}$  and  $2\hat{i} + \hat{j} - 2\hat{k}$  is

- |                     |                      |
|---------------------|----------------------|
| (A) $\frac{\pi}{2}$ | (B) $\frac{\pi}{4}$  |
| (C) $\frac{\pi}{6}$ | (D) $\frac{3\pi}{4}$ |

**Sol.** (B), (D)

Vector  $\vec{AB}$  is parallel to  $[(2\hat{i} + 3\hat{k}) \times (4\hat{j} - 3\hat{k})] \times [(\hat{j} - \hat{k}) \times (3\hat{i} + 3\hat{j})] = 54(\hat{j} - \hat{k})$

Let  $\theta$  is the angle between the vector, then

$$\cos \theta = \pm \left( \frac{54 + 108}{3.54\sqrt{2}} \right) = \pm \frac{1}{\sqrt{2}}$$

$$\text{Hence } \theta = \frac{\pi}{4}, \frac{3\pi}{4}.$$

20.  $f(x) = \begin{cases} e^x, & 0 \leq x \leq 1 \\ 2 - e^{x-1}, & 1 < x \leq 2 \\ x - e, & 2 < x \leq 3 \end{cases}$  and  $g(x) = \int_0^x f(t) dt$ ,  $x \in [1, 3]$  then  $g(x)$  has

- (A) local maxima at  $x = 1 + \ln 2$  and local minima at  $x = e$
- (B) local maxima at  $x = 1$  and local minima at  $x = 2$
- (C) no local maxima
- (D) no local minima

**Sol.** (A), (B)

$$g'(x) = f(x) = \begin{cases} e^x, & 0 \leq x \leq 1 \\ 2 - e^{x-1}, & 1 < x \leq 2 \\ x - e, & 2 < x \leq 3 \end{cases}$$

$g'(x) = 0$ , when  $x = 1 + \ln 2$  and  $x = e$

$$g''(x) = \begin{cases} -e^{x-1}, & 1 < x \leq 2 \\ 1, & 2 < x \leq 3 \end{cases}$$

$g''(1 + \ln 2) = -e^{\ln 2} < 0$  hence at  $x = 1 + \ln 2$ ,  $g(x)$  has a local maximum  
 $g''(c) = 1 > 0$  hence at  $x = c$ ,  $g(x)$  has local minimum.  
 $\therefore f(x)$  is discontinuous at  $x = 1$ , then we get local maxima at  $x = 1$  and local minima at  $x = 2$ .

### **Section - C**

## Comprehension I

There are  $n$  urns each containing  $n + 1$  balls such that the  $i$ th urn contains  $i$  white balls and  $(n + 1 - i)$  red balls. Let  $u_i$  be the event of selecting  $i$ th urn,  $i = 1, 2, 3, \dots, n$  and  $w$  denotes the event of getting a white ball.

21. If  $P(u_i) \propto i$ , where  $i = 1, 2, 3, \dots, n$ , then  $\lim_{n \rightarrow \infty} P(w)$  is equal to



**Sol.** (B)

$$P(u_i) = k_i$$

$$\sum P(u_i) = 1$$

— 1 —

$$\Rightarrow k = \frac{1}{n(n+1)}$$

$$\lim_{n \rightarrow \infty} P(w) = \lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{2i^2}{n(n+1)^2} = \lim_{n \rightarrow \infty} \frac{2n(n+1)(2n+1)}{n(n+1)^2 6} = \frac{2}{3}$$

22. If  $P(u_i) = c$ , where  $c$  is a constant then  $P(u_n/w)$  is equal to

- (A)  $\frac{2}{n+1}$       (B)  $\frac{1}{n+1}$   
 (C)  $\frac{n}{n+1}$       (D)  $\frac{1}{2}$

**Sol.** (A)

(A)

$$P\left(\frac{u_n}{w}\right) = \frac{c\left(\frac{n}{n+1}\right)}{c\left(\frac{\sum i}{n+1}\right)} = \frac{2}{n+1}.$$

23. If  $n$  is even and  $E$  denotes the event of choosing even numbered urn ( $P(u_i) = \frac{1}{n}$ ), then the value of  $P(w/E)$  is

- (A)  $\frac{n+2}{2n+1}$       (B)  $\frac{n+2}{2(n+1)}$   
 (C)  $\frac{n}{n+1}$       (D)  $\frac{1}{n+1}$

**Sol.** (B)

(B)

$$P\left(\frac{W}{E}\right) = \frac{2+4+6+\dots+n}{\frac{n(n+1)}{2}} = \frac{n+2}{2(n+1)}$$

**Comprehension II**

Suppose we define the definite integral using the following formula  $\int_a^b f(x) dx = \frac{b-a}{2} (f(a) + f(b))$ , for more accurate result for  $c \in (a, b)$   $F(c) = \frac{c-a}{2}(f(a) + f(c)) + \frac{b-c}{2}(f(b) + f(c))$ . When  $c = \frac{a+b}{2}$ ,  $\int_a^b f(x) dx = \frac{b-a}{4}(f(a) + f(b) + 2f(c))$ .

24.  $\int_0^{\pi/2} \sin x dx$  is equal to

- (A)  $\frac{\pi}{8}(1+\sqrt{2})$       (B)  $\frac{\pi}{4}(1+\sqrt{2})$   
 (C)  $\frac{\pi}{8\sqrt{2}}$       (D)  $\frac{\pi}{4\sqrt{2}}$

Sol. (A)

$$\int_0^{\pi/2} \sin x dx = \frac{\frac{\pi}{2}+0}{4} \left( \sin(0) + \sin\left(\frac{\pi}{2}\right) + 2\sin\left(\frac{0+\frac{\pi}{2}}{2}\right) \right)$$

$$= \frac{\pi}{8}(1+\sqrt{2}).$$

25. Data could not be retrieved.

26. If  $f''(x) < 0 \forall x \in (a, b)$  and  $c$  is a point such that  $a < c < b$ , and  $(c, f(c))$  is the point lying on the curve for which  $F(c)$  is maximum, then  $f(c)$  is equal to

- (A)  $\frac{f(b)-f(a)}{b-a}$       (B)  $\frac{2(f(b)-f(a))}{b-a}$   
 (C)  $\frac{2f(b)-f(a)}{2b-a}$       (D) 0

Sol.

(A)

$$(F'(c)) = (b-a) f'(c) + f(a) - f(b)$$

$$F''(c) = f''(c) (b-a) < 0$$

$$\Rightarrow F'(c) = 0 \Rightarrow f'(c) = \frac{f(b)-f(a)}{b-a}.$$

**Comprehension III**

Let ABCD be a square of side length 2 units.  $C_2$  is the circle through vertices A, B, C, D and  $C_1$  is the circle touching all the sides of the square ABCD. L is a line through A.

27. If P is a point on  $C_1$  and Q in another point on  $C_2$ , then  $\frac{PA^2 + PB^2 + PC^2 + PD^2}{QA^2 + QB^2 + QC^2 + QD^2}$  is equal to  
 (A) 0.75      (B) 1.25  
 (C) 1      (D) 0.5

Sol.

(A)

Let A, B, C and D be the complex numbers  $\sqrt{2}, -\sqrt{2}, \sqrt{2}i$  and  $-\sqrt{2}i$  respectively.

$$\Rightarrow \frac{PA^2 + PB^2 + PC^2 + PD^2}{QA^2 + QB^2 + QC^2 + QD^2} = \frac{|z_1 - \sqrt{2}|^2 + |z_1 + \sqrt{2}|^2 + |z_1 + \sqrt{2}i|^2 + |z_1 - \sqrt{2}i|^2}{|z_2 + \sqrt{2}|^2 + |z_2 - \sqrt{2}|^2 + |z_2 + \sqrt{2}i|^2 + |z_2 - \sqrt{2}i|^2} = \frac{|z_1|^2 + 2}{|z_2|^2 + 2} = \frac{3}{4}.$$


Sol.

(C)

Let C be the centre of the required circle.

Now draw a line parallel to L at a distance of r (radius of C<sub>1</sub>) from it.

Now  $CP_1 = AC \Rightarrow C$  lies on a parabola.



29. A line M through A is drawn parallel to BD. Point S moves such that its distances from the line BD and the vertex A are equal. If locus of S cuts M at  $T_2$  and  $T_3$  and AC at  $T_1$ , then area of  $\Delta T_1 T_2 T_3$  is

  - (A)  $\frac{1}{2}$  sq. units
  - (B)  $\frac{2}{3}$  sq. units
  - (C) 1 sq. unit
  - (D) 2 sq. units

Sol.

9


$$\therefore \Delta G = \sqrt{2}$$

$$\therefore AT_1 = T_1G = \frac{1}{\sqrt{2}} \quad [\text{as } A \text{ is the focus, } T_1 \text{ is}$$

Also  $T_2T_3$  is latus rectum  $\therefore T_2T_3 = 4 \times \frac{1}{\sqrt{2}}$

Also  $T_2T_3$  is latus rectum  $\therefore T_2T_3 = 4 \times \frac{1}{\sqrt{2}}$

$$\therefore \text{Area of } \Delta T_1 T_2 T_3 = \frac{1}{2} \times \frac{1}{\sqrt{2}} \times \frac{4}{\sqrt{2}} = 1.$$



## Comprehension IV

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix}, \text{ if } U_1, U_2 \text{ and } U_3 \text{ are columns matrices satisfying.}$$

$AU_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  $AU_2 = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}$ ,  $AU_3 = \begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$  and  $U$  is  $3 \times 3$  matrix whose columns are  $U_1, U_2, U_3$  then answer the following questions



Sol. (A)

(A)

Let  $\mathbf{U}_1$  be  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$  so that

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

$$\text{Similarly } U_2 = \begin{bmatrix} 2 \\ -1 \\ -4 \end{bmatrix}, U_3 = \begin{bmatrix} 2 \\ -1 \\ -3 \end{bmatrix}.$$

$$\text{Hence } U = \begin{bmatrix} 1 & 2 & 2 \\ -2 & -1 & -1 \\ 1 & -4 & -3 \end{bmatrix} \text{ and } |U| = 3.$$

31. The sum of the elements of  $U^{-1}$  is

- |        |       |
|--------|-------|
| (A) -1 | (B) 0 |
| (C) 1  | (D) 3 |

**Sol.** **(B)**

$$\text{Moreover } \text{adj } U = \begin{bmatrix} -1 & -2 & 0 \\ -7 & -5 & -3 \\ 9 & 6 & 3 \end{bmatrix}.$$

$$\text{Hence } U^{-1} = \frac{\text{adj } U}{3} \text{ and sum of the elements of } U^{-1} = 0.$$

32. The value of  $[3 \ 2 \ 0] U \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}$  is

- |       |         |
|-------|---------|
| (A) 5 | (B) 5/2 |
| (C) 4 | (D) 3/2 |

**Sol.** **(A)**

$$\text{The value of } [3 \ 2 \ 0] U \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}$$

$$= [3 \ 2 \ 0] \begin{bmatrix} 1 & 2 & 2 \\ -2 & -1 & -1 \\ 1 & -4 & -3 \end{bmatrix} \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}$$

$$= [-1 \ 4 \ -4] \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix} = -3 + 8 = 5.$$

### Section - D

33. If roots of the equation  $x^2 - 10cx - 11d = 0$  are  $a, b$  and those of  $x^2 - 10ax - 11b = 0$  are  $c, d$ , then the value of  $a + b + c + d$  is ( $a, b, c$  and  $d$  are distinct numbers)

**Sol.** As  $a + b = 10c$  and  $c + d = 10a$

$$\begin{aligned} ab &= -11d, cd = -11b \\ \Rightarrow ac &= 121 \text{ and } (b + d) = 9(a + c) \\ a^2 &- 10ac - 11d = 0 \\ c^2 &- 10ac - 11b = 0 \\ \Rightarrow a^2 + c^2 - 20ac - 11(b - d) &= 0 \\ \Rightarrow (a + c)^2 - 22(121) - 11 \times 9(a + c) &= 0 \\ \Rightarrow (a + c) &= 121 \text{ or } -22 \text{ (rejected)} \\ \therefore a + b + c + d &= 1210. \end{aligned}$$

34. The value of  $\frac{5050 \int_0^1 (1-x^{50})^{100} dx}{\int_0^1 (1-x^{50})^{101} dx}$  is

$$\begin{aligned} \text{Sol. } &= \frac{5050 \int_0^1 (1-x^{50})^{100} dx}{\int_0^1 (1-x^{50})^{101} dx} = 5050 \frac{I_{100}}{I_{101}} \\ I_{101} &= \int_0^1 (1-x^{50})(1-x^{50})^{100} dx \\ &= I_{100} - \int_0^1 x \cdot x^{49} (1-x^{50})^{100} dx \\ &= I_{100} - \left[ \frac{-x(1-x^{50})^{101}}{101} \Big|_0^1 - \int_0^1 \frac{(1-x^{50})^{101}}{5050} dx \right] \\ I_{101} &= I_{100} - \frac{I_{101}}{5050} \\ \Rightarrow \quad &5050 \frac{I_{100}}{I_{101}} = 5051. \end{aligned}$$

35. If  $a_n = \frac{3}{4} - \left( \frac{3}{4} \right)^2 + \left( \frac{3}{4} \right)^3 + \dots + (-1)^{n-1} \left( \frac{3}{4} \right)^n$  and  $b_n = 1 - a_n$ , then find the minimum natural number  $n_0$  such that  $b_n > a_n \forall n \geq n_0$

$$\begin{aligned} \text{Sol. } a_n &= \frac{3}{4} - \left( \frac{3}{4} \right)^2 + \left( \frac{3}{4} \right)^3 + \dots + (-1)^{n-1} \left( \frac{3}{4} \right)^n \\ &= \frac{\frac{3}{4} \left( 1 - \left( -\frac{3}{4} \right)^n \right)}{1 + \frac{3}{4}} = \frac{3}{7} \left( 1 - \left( -\frac{3}{4} \right)^n \right) \\ b_n > a_n \Rightarrow 2a_n < 1 & \\ \Rightarrow \frac{6}{7} \left( 1 - \left( -\frac{3}{4} \right)^n \right) < 1 & \\ \Rightarrow 1 - \left( -\frac{3}{4} \right)^n < \frac{7}{6} & \\ \Rightarrow -\frac{1}{6} < \left( -\frac{3}{4} \right)^n \Rightarrow \text{minimum natural number } n_0 = 6. & \end{aligned}$$

36. If  $f(x)$  is a twice differentiable function such that  $f(a) = 0, f(b) = 2, f(c) = -1, f(d) = 2, f(e) = 0$ , where  $a < b < c < d < e$ , then the minimum number of zeroes of  $g(x) = (f'(x))^2 + f''(x)f(x)$  in the interval  $[a, e]$  is

$$\text{Sol. } g(x) = \frac{d}{dx}(f(x) \cdot f'(x))$$

to get the zero of  $g(x)$  we take function

$$h(x) = f(x) \cdot f'(x)$$

between any two roots of  $h(x)$  there lies at least one root of  $h'(x) = 0$

$$\begin{aligned} \Rightarrow g(x) = 0 \\ h(x) = 0 \end{aligned}$$

- $\Rightarrow f(x) = 0$  or  $f'(x) = 0$   
 $f(x) = 0$  has 4 minimum solutions  
 $f'(x) = 0$  has 3 solution  
 $h(x) = 0$  has 7 solution  
 $\Rightarrow h'(x) = g(x) = 0$  has minimum 6 solutions.

## Section - E

37.

Match the following:

Normals are drawn at points P, Q and R lying on the parabola  $y^2 = 4x$  which intersect at  $(3, 0)$ . Then

- |  |                |
|--|----------------|
| (i) Area of $\triangle PQR$                    | (A) 2          |
| (ii) Radius of circumcircle of $\triangle PQR$ | (B) $5/2$      |
| (iii) Centroid of $\triangle PQR$              | (C) $(5/2, 0)$ |
| (iv) Circumcentre of $\triangle PQR$           | (D) $(2/3, 0)$ |

Sol.

As normal passes through  $(3, 0)$ 

$$\Rightarrow 0 = 3m - 2m - m^3$$

$$\Rightarrow m^3 - m \Rightarrow m = 0, \pm 1$$

$$\therefore \text{Centroid} = \left( \frac{(m_1^2 + m_2^2 + m_3^2)}{3}, -\frac{2(m_1 + m_2 + m_3)}{3} \right) = \left( \frac{2}{3}, 0 \right)$$

$$\text{Circumcentre (mid point of PR)} = \left( \frac{m_1^2 + m_2^2}{2}, -(m_1 + m_2) \right) = (1, 0).$$

$$\text{Circum radius} = \sqrt{\frac{-2m_1 + 2m_2}{2}} = 2 \text{ units.}$$

$$Q \equiv (m_2^2, -2m_2) \equiv (1, -2)$$

$$R \equiv (m_3^2, -2m_3) \equiv (1, 2)$$

$$\text{Area of } \triangle PQR = \frac{1}{2} \times 4 \times 1 = 2 \text{ sq. units.}$$

$$R = \frac{QR}{2 \sin \angle QPR} = \frac{4}{\sin(2 \tan^{-1} 2)}$$

$$\Rightarrow \frac{4}{2 \times \sin \left( \tan^{-1} \frac{4}{1-4} \right)} = \frac{4}{2 \times \frac{4}{5}} = \frac{5}{2}$$

$$\therefore \text{circumcentre} \equiv \left( \frac{5}{2}, 0 \right).$$

38.

Match the following

- |   |               |
|---|---------------|
| (i) $\int_0^{\pi/2} (\sin x)^{\cos x} (\cos x \cot x - \log(\sin x)^{\sin x}) dx$             | (A) 1         |
| (ii) Area bounded by $-4y^2 = x$ and $x - 1 = -5y^2$  | (B) 0         |
| (iii) Cosine of the angle of intersection of curves $y = 3^{x-1} \log x$ and $y = x^x - 1$ is | (C) $6 \ln 2$ |
| (iv) Data could not be retrieved.   | (D) $4/3$     |

$$\text{Sol. (i)} \quad I = \int_0^{\pi/2} (\sin x)^{\cos x} (\cos x \cdot \cot x - \log(\sin x)^{\sin x}) dx$$

$$\Rightarrow I = \int_0^{\pi/2} \frac{d}{dx} (\sin x)^{\cos x} dx = 1.$$

- (ii) The points of intersection of  $-4y^2 - x$  and  $x - 1 - 5y^2$  is  $(-4, -1)$  and  $(-4, 1)$

$$\text{Hence required area} = 2 \left[ \int_0^1 (1 - 5y^2) dy - \int_0^1 -4y^2 dy \right] = \frac{4}{3}.$$

- (iii) The point of intersection of  $y = 3^{x-1} \log x$  and  $y = x^x - 1$  is  $(1, 0)$

$$\text{Hence } \frac{dy}{dx} = \frac{3^{x-1}}{x} + 3^{x-1} \log 3 \cdot \log x. \quad \left. \frac{dy}{dx} \right|_{(1,0)} = 1$$

$$\text{for } y = x^x - 1. \quad \left. \frac{dy}{dx} \right|_{(1,0)} = 1$$

If  $\theta$  is the angle between the curve then  $\tan \theta = 0 \Rightarrow \cos \theta = 1$ .

$$(iv) \frac{dy}{dx} = \left( \frac{2}{x+y} \right)$$

$$\Rightarrow \frac{dx}{dy} = \frac{x}{2}$$

$$\Rightarrow xe^{-y/2} = \frac{1}{2} \int y \cdot e^{-y/2} dy$$

$$\Rightarrow x + y + 2 = ke^{y/2} = 3e^{y/2}.$$

39.

Match the following

- (i) Two rays in the first quadrant  $x + y = |a|$  and  $ax - y = 1$  intersects each other in the interval  $a \in (a_0, \infty)$ , the value of  $a_0$  is

(A) 2

- (ii) Point  $(\alpha, \beta, \gamma)$  lies on the plane  $x + y + z = 2$ . Let

(B)  $4/3$

$$\bar{a} = \alpha \hat{i} + \beta \hat{j} + \gamma \hat{k}, \quad \hat{k} \times (\hat{k} \times \bar{a}) = 0, \text{ then } \gamma = .$$

$$(iii) \left| \int_0^1 (1-y^2) dy \right| + \left| \int_1^0 (y^2-1) dy \right|$$

(C)  $\left| \int_0^1 \sqrt{1-x} dx \right| + \left| \int_{-1}^0 \sqrt{1+x} dx \right|$

- (iv) If  $\sin A \sin B \sin C + \cos A \cos B = 1$ , then the value of  $\sin C =$

(D) 1

**Sol.** (i) Solving the two equations of ray i.e.  $x + y = |a|$  and  $ax - y = 1$

$$\text{we get } x = \frac{|a|+1}{a+1} > 0 \text{ and } y = \frac{|a|-1}{a+1} > 0$$

when  $a+1 > 0$ ; we get  $a > 1 \therefore a_0 = 1$ .

- (ii) We have  $\bar{a} = \alpha \hat{i} + \beta \hat{j} + \gamma \hat{k} \Rightarrow \bar{a} \cdot \hat{k} = \gamma$

$$\text{Now; } \hat{k} \times (\hat{k} \times \bar{a}) = (\hat{k} \cdot \bar{a}) \hat{k} - (\hat{k} \cdot \hat{k}) \bar{a}$$

$$= \gamma \hat{k} - (\alpha \hat{i} + \beta \hat{j} + \gamma \hat{k})$$

$$= \alpha \hat{i} + \beta \hat{j} = \vec{0} \Rightarrow \alpha = \beta = 0$$

As  $\alpha + \beta + \gamma = 2 \Rightarrow \gamma = 2$ .

$$(iii) \left| \int_0^1 (1-y^2) dy \right| + \left| \int_1^0 (y^2-1) dy \right|$$

$$= 2 \int_0^1 (1-y^2) dy = \frac{4}{3}$$

$$\left| \int_0^1 \sqrt{1-x} dx \right| + \left| \int_{-1}^0 \sqrt{1+x} dx \right| = 2 \int_0^1 \sqrt{1-x} dx$$

$$= 2 \int_0^1 \sqrt{x} dx = 2 \cdot \frac{2}{3} \cdot x^{3/2} \Big|_0^1 = \frac{4}{3}.$$

- (iv)  $\sin A \sin B \sin C + \cos A \cos B \leq \sin A \sin B + \cos A \cos B - \cos(A - B)$   
 $\Rightarrow \cos(A - B) \geq 1 \Rightarrow \cos(A - B) = 1 \Rightarrow \sin C = 1.$

40. Match the following

(i)  $\sum_{i=1}^{\infty} \tan^{-1} \left( \frac{1}{2i^2} \right) = t$ , then  $\tan t =$  (A) 0

(ii) Sides  $a, b, c$  of a triangle  $\Delta ABC$  are in AP and

$$\cos \theta_1 = \frac{a}{b+c}, \cos \theta_2 = \frac{b}{a+c}, \cos \theta_3 = \frac{c}{a+b}, \text{ then } \tan^2 \left( \frac{\theta_1}{2} \right) + \tan^2 \left( \frac{\theta_3}{2} \right) = \text{(B) 1}$$

(iii) A line is perpendicular to  $x + 2y + 2z = 0$  and passes through  $(0, 1, 0)$ . (C)  $\frac{\sqrt{5}}{3}$

The perpendicular distance of this line from the origin is

(D)  $2/3$

(iv) Data could not be retrieved.

Sol. (i)  $\sum_{i=1}^{\infty} \tan^{-1} \left[ \frac{1}{2i^2} \right] = t$

$$\begin{aligned} \text{Now; } & \sum_{i=1}^{\infty} \tan^{-1} \left[ \frac{2}{4i^2 - 1 + 1} \right] \\ &= \sum_{i=1}^{\infty} \left[ \tan^{-1}(2i+1) - \tan^{-1}(2i-1) \right] \\ &= \left[ (\tan^{-1} 3 - \tan^{-1} 1) + (\tan^{-1} 5 - \tan^{-1} 3) + \dots + \tan^{-1}(2n+1) - \tan^{-1}(2n-1), \dots, \infty \right] \end{aligned}$$

$$t = \tan^{-1}(2n+1) - \tan^{-1} 1 = \tan^{-1} \frac{2n}{1+(2n+1)}$$

$$\Rightarrow \tan t = \frac{n}{n+1} \Rightarrow t = \frac{\pi}{4}$$

(ii) We have  $\cos \theta_1 = \frac{1 - \tan^2 \frac{\theta_1}{2}}{1 + \tan^2 \frac{\theta_1}{2}} = \frac{a}{b+c} \Rightarrow \tan^2 \frac{\theta_1}{2} = \frac{b+c-a}{b+c+a}$

$$\text{Also, } \cos \theta_3 = \frac{1 - \tan^2 \frac{\theta_3}{2}}{1 + \tan^2 \frac{\theta_3}{2}} = \frac{c}{a+b} \Rightarrow \tan^2 \frac{\theta_3}{2} = \frac{a+b-c}{a+b+c}$$

$$\therefore \tan^2 \frac{\theta_1}{2} + \tan^2 \frac{\theta_3}{2} = \frac{2b}{3b} = \frac{2}{3}$$

(iii) Line through  $(0, 1, 0)$  and perpendicular to plane  $x + 2y + 2z = 0$  is given by  $\frac{x-0}{1} = \frac{y-1}{2} = \frac{z-0}{2} = r$ .

Let  $P(r, 2r+1, 2r)$  be the foot of perpendicular on the straight line then

$$r \times 1 - (2r+1) 2 + 2 \times 2r = 0 \Rightarrow r = -\frac{2}{9}$$

$$\therefore \text{Point is given by } \left( -\frac{2}{9}, \frac{5}{9}, -\frac{4}{9} \right)$$

$$\therefore \text{Required perpendicular distance} = \sqrt{\frac{4+25+16}{81}} = \frac{\sqrt{5}}{3} \text{ units.}$$

(iv) Data could not be retrieved.