
JIPMER 2001

1)Vocalis Muscle is a part of
a)Thyroarytenoid
b)Cricoarytenoid
c)Transverse arytenoid
d)Posterior cricoarytenoid
2)Sumatriptan is contraindicated in
a)Asthma
b)DM
c)Coronary artery disease
d)peripheral vascular disease
3)Non granular fungal disease is
a)Cryptococcus
b)Blastomycosis
c)Candida
d)
4)All following are true of seminiferous tubules except
a)Sertoli cells
b)Leydig cells
c)spermatid

d)Spermatogonia
5)Sub-poena is a
a)Legal
b)medicolegal
c)Medical
d)first info report
6)Refsum disease is due to defect in enzyme
a)Phytanic acid oxidase
b)
c)
d)
7)Skin supplied over the angle of mandible and parotid area is by
a)Greater auricular
b)posterior auricular
c)auriculotemporal
d)facial
8)Adductor pollicis is supplied by
a)Superficial branch of ulnar nerve
b)deep branch of ulnar nerve

c)radial nerve d)median nerve
9)Ring Scotoma is seen in
a)Retinitis Pigmentosa
b)
c)
d)
10) In a patient, highest visual morbidity is seen in
a)Nuclear cataract
b)Intumescent cataract
c)Posterior subcapsular
d)
11)Klumpke paralysis is
a)C5,C6
B)C6,C7
C)C8,T1
D)T1,T2
12)For the treatment of a patient with after cataract which laser is used
a)LASIC
b)YAG

c)excimer
d)Argon
13)Most malignant breast tumor of the following is
a)inflammatory carcinoma
b)Mucinous carcinoma
c)Colloid carcinoma
d)Paget's` disease of breast.
14)Full thickness graft means:
a)Only epidermis
b)epidermis and superficial dermis
c)epidermis + whole dermis
d)epidermis, dermis and subcutaneous tissue
15)A 50yr old woman weighing 50kg and having almost complete burns comes to burns ward. According to Parklands formula ,the amt. of fluid for resuscitation in 24hrs.
a)5L
b)10L
c)12L
d)14L
16)Investigation of choice for DVT is
a)Duplex



a)Giant cell tumor
b)osteoclastoma
c)
d)
21)Snow storm appearance on CXR is seen in
a)anthracosis
b)asbestosis
c)Silicosis
d)Bagassosis
22)Housemaid Knee is inflammation of
22)Housemaid Knee is inflammation of a)prepatellar bursa
a)prepatellar bursa
a)prepatellar bursa b)Suprapatellar bursa
a)prepatellar bursa b)Suprapatellar bursa c)Infrapatellar bursa
a)prepatellar bursa b)Suprapatellar bursa c)Infrapatellar bursa
a)prepatellar bursa b)Suprapatellar bursa c)Infrapatellar bursa d)anserine bursa
a)prepatellar bursa b)Suprapatellar bursa c)Infrapatellar bursa d)anserine bursa 23) Height increase in adolescent/year is
a)prepatellar bursa b)Suprapatellar bursa c)Infrapatellar bursa d)anserine bursa 23) Height increase in adolescent/year is a)8.5 cm

24)Patient with genotype XO will have following phenotype except
a)Tall stature
b)Broad Chest
c)Webbed neck
d)Lymphedema
25)treatment of carcinoma thyroid using radioactive iodine is
a) I131
b) I125
c) I133
26) solitary nodule of thyroid in a male aged 40yr from a non-endemic area would most likely have
a)MNG
b)papillary ca Thyroid
c)follicular ca
d)
27)In assessment of gestational age of a neonate the following all seen except
a)Breast bud
b)Ocular
c)Genitalia
d)scrotal skin

28)On USG, the appendix is seen, it is most likely due to
a)Retrocecal appendix
b)Acute appendicitis
d)preileal appendix
29)Vesicular lesion is seen in
a)Primary syphilis
b)secondary syphilis
c)tertiary syphilis
d)congenital syphilis
30)In villous adenoma of rectum which of the following is seen
a) Hypokalemia
b)Hyperkalemia
c)hypochloremic alkalosis
d)
31)The best view on X-ray to view pituitary fossa is
a)lateral skull X-ray
b)Towne's view
c)schuller's view
d)

32)Supratendinitis calcification is due to deposition of
a)Ca2+ pyrophosphate
b) Ca2+ hydroxyapatite
c)Ca2+ oxalate
d)
33) all the following are beta-blockers except
a)Esmolol
b)sotalol
c) celiprolol
d)bambuterol
34)Unpleasant dreams and hallucinations occur after which anesthesia
a)Ketamine
b)Halothane
c)thiopentone
d)
35)Longest acting antihistamine is
a)Cetrizine
b)Astemizole
c)terfenadine
d)acrivastive

36) The most common carcinoma of sinuses is
a) squamous cell carcinoma
b) adeno carcinoma
c) transitional cell carcinoma
d)
38) All are specific to TB except
a) Epituberculosis
b) Phlycten
c) Erythema nodosum
d) lupus vulgaris
39) The following macrolides are used for toxoplasmosis except
39) The following macrolides are used for toxoplasmosis except a) Erythromycin
a) Erythromycin
a) Erythromycin b) clarithromycin
a) Erythromycin b) clarithromycin c) Azithromycin
a) Erythromycin b) clarithromycin c) Azithromycin
a) Erythromycin b) clarithromycin c) Azithromycin d) spiramycin
 a) Erythromycin b) clarithromycin c) Azithromycin d) spiramycin 40) Following are third generation cephalosporins except

d) cefotaxime
41) The mechanism of action of cephalosporin is
a) interferes with cell wall synthesis
b) inhibition of DNA gyrase
c)
d)
42) Evidence of ovulation has occurred is all except
a) 10 mm graffian follicles on USG**
b) progesterone > 5ng/ml
c) corpus luteum
d) Biphasic increase in temperature
43) Objective evidence of hearing loss is all except
a) pure tone audiometry
b) BAER
c) otoacoustic emission
d)
44) Vestibular anus is
a) imperforate anus
b) high anorectal

b) a state motor or verbal restlessness
c) cogwheel rigidity
d)
49)infantile autism is seen in
a) dysthymic personality
b) personality disorder
b) paranoid personality
d) behavioral disorder
50) In pulmonary embolism, the receptor involving causing reflex hyperventilation is
a) J-receptor
b) slow adapting receptor
c) fast adapting receptor
d) Irritant receptors
51) Cazt** channel defect is seen in
a) hypokalemic periodic paralysis
b) hyperkalemic periodic paralysis
c) paramyotonia congenita
d)
52) Which is not an aldolase

56) A patient comes with recurrent abdominal pain, and urinary examination reveals Ehrlich aldehyde test (+) for urobilonogens. The diagnosis is
a) porphyria
b) renal calculi
c) cholelithiasis
d)
57) In urticaria pigmentosa, the cell most commonly seen in skin on microscopy is
a) Mast cell
b) Langerhan's cell
c) Langhan's cell
d)
58)lagopathalmos is seen in
a) leprosy
b) syphilis
c) peripheral neuritis
d) HZV infection
59) Topical steroids is used in
a) Interstitial keratitis
b) bacterial keratitis
c) pendrotic
d) herpetic

60) An elderly male with dry skin and fissuring scaly skin is likely to have
a) nummular eczema
b) venous eczema
c) asteatotic eczema
d)
61) In myasthenia gravis, true is
a) antibody is present against nicotinic receptor
b) decreased Acetylcholine release
c) decreased no of receptors
d) end organ resistance
62) Insulin sensitivity increases on following treat except
a) metformin
b) acarbose
c) exercise
d) fasting
63) All the following are anticonvulsants except
a) lamotrigine
b) methyl phenytoline
c) vigabatrin

d) topiramate
64) Side effect of valvular heart disease is seen with
a) fenfluramine
b) methylphenidate
c) amphetamine
d)
65) Side effect of acute pancreatitis is due to following anti HIV drugs
a) lamivudine
b) zidovudine
c) didanosine
d)
66) "milk spots" on the omentum consists of
a) macrophage
b) neutrophils
c) lymphocyte
d)
67) longest acting muscle relaxant is
a) pancuronium
b) mivacurium

c) vecuronium
d)
68)Genital prolapse in a female can be due to all except
a) Ascites
b)complete perineal tear
c) chronic cough
d)
69) Tubal patency test is done by which test?
a) sion test
b) frei's test
c) bonney's test
d) three swab test
70) cells are most susceptible to radiation in which phase
a) M phase
b) G2 phase
c) G1 phase
d) S phase
Ans. Ref:
71) end product of ?- hydroxylation is

a) Acetyl CoA + propionyl CoA
b) Acetyl CoA + succinyl CoA
c) Acetyl CoA +
d) Acetyl CoA + malonyl CoA
72) Morrison pouch is
a) subphrenic space
b) posterior to lateral lobe of liver
c) lesser sac
73) hemoptysis in mitral stenosis is due to
a) Pulmonary arterial hypertension
b) Pulmonary arterial hypertension + left atrial hypertension
c) Left ventricular pressure
d)
74) In massive pericardial effusion all are seen except a) on fluoroscopy less prominent vascular marking
b) CT ratio >0.5
c) CP angles acute
d) Plethoric pulmonary fields
75) single plate coated with ammonium**** is present in all except
a) X-ray

b) MMR
c) CT scan
d) USG film
76)Best view on x-ray to view bilateral auditory oscines /Eustachian tubes? is
a) occipito frontal
b) fronto occipital
c) submento vertical
d)
77) best view for pituitary fossa on x-ray is
a) lateral skull x-ray
b)
c) sella view
d)
78) Gram negative antigen is injected in a rabbit after 2 weeks****question not complete***
a) schwartzmann reaction- ans
b) Schultz-dale phenomenon
c) P-K-reaction
d)
79) The most common cause of spasm of psoas in childhood is

a) trauma
b) tuberculosis psoas
c) acute appendicitis
d) pyogenic psoas abscess
80) which leucotriene is the adhesion factor for the neutrophil on the cell surface to attach endothelium
a) B4
b) C4
c)D4
d) E4
81) In the following conditions, bone matrix is reduced except
a) osteoporosis
b) osteomalacia
c) hyperparathyroidism
d) osteopetrosis
discuss tips or queries at message board(a new window will open)
82) which one is FAD dependent reaction
a) succinyl CoA fumarate
b)
c) d)

83) which of the following is formed by the intestinal mucosal cells on absorption of dietary lipids
a) chylomicrons
b) VLDL
c) HDL
d) LDL
84) high risk factor for coronary artery diseases are all except
a) low homocysteine level
b) cholesterol > 250 mg
c)
d)
85) In osteomalacia, all the following seen except
a) looser zones
b) high alkaline phosphate levels
c)
d)
86) In a patient with solitary nodule of thyroid, investigation of choice is
a) FNAC
b) USG
c) CT
d) Radioisotope scan

87) how many double bonds are present in arachidonic acid?
a) One
b) two
c) Three
d) Four
88) In elective tracheostomy, incision is taken on
a) cricothyroid membrane
b) C1 & C2
c) c2 & c4
d)
89) Autoinduction of biotransformation and action as a hormone is seen in
89) Autoinduction of biotransformation and action as a hormone is seen in a) carbamazepine
a) carbamazepine
a) carbamazepineb) cimetidine
a) carbamazepineb) cimetidine
a) carbamazepineb) cimetidinec) corticosteroid.
a) carbamazepineb) cimetidinec) corticosteroid.90) G-6-P-D inheritance is
 a) carbamazepine b) cimetidine c) corticosteroid. 90) G-6-P-D inheritance is a) X-linked - recessive

91) oral hairy leucoplakia & whitish plaque on the tongue and palate seen gives clue to most likely diagnosis of
a) HIV infection
b) HTLV - 1
c) infectious mononucleosis
d)
92) most common organism causing spontaneous bacterial peritonitis is
a) staphylococcus
b)
c) salmonella
d) streptococcus
93) symblepharon is caused by
a) diphtheria
b) gonococci
c) chlamydia
d) opthalmia neonatorum
94) jargon aphasia is seen in
a) wenicke's
b) broca's aphasia
c) conduction aphasia

95) In koraskoff psychosis seen is
a) global deterioration intelligence
b) impaired recall & new ability to learn
c)
d)
96) multiple calcification in the brain or cystic calcification in the brain or comma shaped calcification seen in
a) CMV
b) neurocysticercosis
c) tuberous sclerosis
d) toxoplasmosis
97)On CT scan, all are seen as hypodense area except
a) cerebral hemorrhage
b) glioblastoma
c) cerebral edema
d) cerebral infarct
98) hypertensive bleed mostly occurs in
a)putamen
b) pons
c) cortex

d) thalamus
99) back-wash ileitis is seen in
a) crohn's diseases
b) ulcerative colitis
c)
d)
100) leukemoid reaction type 2 in leprosy is due to
a) type I hypersensitivity
b) Type II hypersensitivity
c)type III hypersensitivity
d) type IV hypersensitivity
101) systematic desensitization as form of therapy is done in
a) phobia
b) obsessive compulsive disorder
c)
d)
102) morbid jealousy is seen in
a) alcoholism
b) persons who are isolated from society

c)
d)
103) respiratory alkalosis is seen in
a) duodenal obstruction
b) early salicylism
c) diarrhea
d)
104) dopamine is highest in
a)striatum
b) cerebellum
c) caudate
d)
105) IL-1 & TNF ? causes
a) increasing capillary permeability
c) acute phase reactants
b) chemotaxis
d) activate endothelium
106) thimble pitting nails are seen in
a) psoriasis

b) tinea versicolor
c)lichen planus
d)tuberculosis
107) Ridley-Jopling classification includes all except
a) lepromatous leprosy
b) borderline borderline
c) tuberculoid
d) neuritic
108) which of the following is type IV hyper sensitivity
a)
b)
c)
d) contact dermatitis
109) Pressure necrosis which causes venous flow to stop during endotracheal intubation occurs at
a) 20 - 30 mm Hg
b) 15 - 20 mm Hg
c) 30-40 mm Hg
d) 16 -15 mm Hg
110) diagnosis of mucopolysacharridoses can be done by examination of

a) cornea
b) skin
c) bone
d) face
111) cadaveric position of vocal cords occurs in
a) recurrent laryngeal nerve palsy
b) recurrent laryngeal nerve palsy + superior laryngeal nerve palsy
c)
d)
112) most common site of pelvic abscess is in
a) ishiorectal fossa
b) perianal
c)
d)
113) most important sensory organ in the ampulla is
a) organ of corti
b) macula
c)otolithi
d) crista

114) carotid body baroreceptor is most sensitive to
a)mean blood pressure
b) diastolic blood pressure
c) systolic blood pressure
d) Pulse pressure
115) the following are derived from endoderm except
a) liver
b) kidney
c) pancreas
d)urinary bladder
116)human placenta is
a) hemochorial
b) endothen - esdothelima
c) chorio - endothelial
d)
117)In malaria acquired by transfusion, the infective form is
a) sporozoites
b) shistozoites
c) gametocytes
d) merozoites

118) inhibition of prolactin is caused by
a) dopamine
b) dobutamine
c)
d)
119) In after coming, head best forceps used for delivery is
a) piper's forceps
b) kielland's forceps
c) wrigley' forceps
d)
120) longest diameter of the skull is
a) suboccipital bregmatic
b) occipital frontal
c) bitemporal
d) biparietal
121)bishop score is used to evaluate
a) cervix status
b) uterine contraction
c) fetal well being

d)
122) the management of eclampsia includes all except
a) control of convulsion
b) prolongation of pregnancy
c) control of BP
d) delivery of fetus
123) The most common cause of vesicovaginal fistula in developing countries is
a) pressure necrosis of urethra during obstructed labor
b) forceps delivery
c)
d)
124) shoulder dystocia occurs in delivery of fetus with
a) anencephaly
b) hydrocephalus
c) fetal ascites
d)
125) benign prost hyperplasia occur in the
a) periurethral transitional zone
b) cortex

c) posterior lobe
d)
126) tetany is seen in all except
a) hypoparathyroidism
b) full strength milk given in neonate
c)
d)
127) benign intracranial hypertension is seen in all except
a) hypertvitaminosis A
b) steroid therapy
c) OCP
d)
128) the most common breech presentation in a nulliparous woman is
a) frank breech
b) complete breech
c) footling
d)
129) In supra condylar fracture, the x-ray view of which helps in the next line of management
a) baumman's angle

b) cobb's angle
c) cowen's angle
d)
discuss tips or queries at AIPPG message board(a new window will open)
130) calculate the pearl index if 6 pregnancy occurs in 1200 females practicing contraception for 1 year
a) 6
b) 12
c) 18
d) 24
131) In minilap procedure, the most common method successfully used is
a) uchida
b) pomeroy's
c)irving's
d)
132) deep transverse arrest occurs in
a) android
b) anthropoid
c) gynecoid
d) flat pelvis
133) false positive rinne's test occur in

a) unilateral conductive deafness
b) unilateral sensory neural deafness
c) bilateral conductive deafness
d) bilateral sensory neural deafness
134) A nulliparous female aged 30 years develops IInd degree UV prolapse. The operative procedure done is
a) cervicopexy
b)LeForte
c)Manchester
135) following are true of hemorrhoids except
a) they are arteriolar dilatations
b) they are common causes of painless bleeding
c) they cannot be per rectally palpated
136) waves on EEG during NREM period of sleep is
a) alpha
b) beta
c) delta
d) theta
137) following is true of hormone leptin except
a) it is found in increased levels in obese

b) it's site of action is hypothalmus
c) it is produced by adipose all
d) it increases appetite.
138) function of brown fat is
a) glycogenosis
b) gluconeogenesis
c)
d) thermogenesis
139) The most common cause for stridor in the early day of life is
a) laryngomalacia
b)esophago- tracheal fistula
c) laryngeal stenosis
d)
140) commonest sources of lead to cause increased blood level is from
a) air
b) lead paints
c) water
d)
141) The incisura of arterial pulse corresponds with

a)S1
b) S2
c) opening snap
4) S3
142) the one organism which is responsible for coronary artery disease
a) chlamydia
b) klebsiella
c) E coli
d) mycoplasma
143) to build a genome library using digested chromosome into fragments, the enzyme used is
a) DNA polymerase
b) RNA restriction endonuclease
d)
144) all the following are suggestive of urethral injury except
a) blood at meatus
b) haematoma of scrotum
c) high prostate on PR
d) blood in rectum
145) to assess the refraction state of eye, the investigation done is

a) USG - Bscan
b) retinoscopy
c) binoculoscopy
d) opthalmoscopy
146) which of the following has the worst prognosis
a) rolandic epilepsy
b) versive epilepsy
c) absence epilepsy
d)
Ans. Ref:
147) which of the following has worst prognoses in carcinoma lung
a) squamous cell carcinoma
b) large cell carcinoma
c) small cell carcinoma
d)adenocarcinoma
148) Herbert's pit is seen in
a) vernal conjunctivitis
b) atopic conjunctivitis
c) gonococcal conjunctivitis
d)chlamydial conjunctivitis

149) the amount of rads required for treatment of oropharyngeal carcinoma is
a) 3000 - 4000 rads
b) 4000- 5000 rads
c) 5000 - 6000
d) 6000 - 7000
150) prolonged ST segment with T wave seen in
a) hypokalemia
b) hyperkalemia
c) hypomagnesemia
d) hypocalcemia
151) In a meconium aspirated baby following are investigations. pO2=50, pCO2=60 & blood pH=7.1. the next line of management of giving I V infusion to correct this condition is
pH=7.1. the next line of management of giving I V infusion to correct this condition is
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate c) CPAP
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate c) CPAP
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate c) CPAP d) endotracheal intubation & IPPV
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate c) CPAP d) endotracheal intubation & IPPV 152) which of the following hormones increase during sleep
pH=7.1. the next line of management of giving I V infusion to correct this condition is a) NAHCO3 solution b) ringer lactate c) CPAP d) endotracheal intubation & IPPV 152) which of the following hormones increase during sleep a) insulin

153) bromocriptine is used to suppress prolactin
a) acts as agonist to dopamine receptors
b)
c)
d)
154) in chronic folliculitis the treatment of choice is
a)
b)
c) cloxacillin
d) vancomycin
Ans. Ref:
155) on X-ray calcification of breast is seen. Likely cause is
a) chronic breast abscess
b) tuberculosis of breast
b) tuberculosis of breastc) periductal carcinoma
c) periductal carcinoma
c) periductal carcinoma
c) periductal carcinoma d)
c) periductal carcinomad)156) epidermolysis bullosa is due to

d) genetic fragility of skin
157) A child suffering from marasmus was given high protein diet. The metabolic change likely is
a) hyperargininemia
b) hyperammonemia
c) hyperlysinemia
d) hypernatremia.
158) the ORS fluid used for a marasmic child lacks in (racemol)
a) amino acids
b) sodium
c) copper
d) magnesium
159) which of the following cell has both T lymphocyte and B lymphocyte function
a) plasma cell
b) T- lymphoblast cell
c)B-lymphoblast cell
d) NK cell
160) salmonella osteomyelitis occurs in
a) sickle cell d
b) beta thalassemia

c) sideroblastic anemia
d)
161) following are louse borne disease
a) endemic typhus
b) KFD
c) lyme's disease
d)
162) the treatment of choice in angiofibroma is a 12 year old boy is
a) radio treatment
b) chemo treatment
c) embolisation of inferior palatine artery
d) surgery
discuss tips or queries at AIPPG message board(a new window will open)
163) A young women, 30 yrs old complains of pain in the left knee joint which increases after exercise since 4 months. An x-ray shows eccentric bone cyst. The likely diagnosis is
a) aneurysmal bone cyst
b) unicameral bone cyst
c)
d)
164) The term family size in demographic parlance means
a) the total number of children a women has borne at a point of time

b) the total number of children of couples
c) the total number of children by women borne in a female between 15-45 yrs
Ans. Ref:
165) the approximate magnitude of completed family size from demographic point view is given by
a) TFR
b) GFR
c)NRR
166) the following should be done for prophylaxis of rabies in a 1 year old child except
a) wash the wound with soap & water
b) give HDCV in the buttock
c) HDCV should not be given to HIV pt
d)
167) all the following are examples of health promotion except
a) stop smoking
b) regular exercise
c) chemoprophylaxis
168) one of the following helps deliver baby in a community
a) health assistant female
b) anganwadi worker
c) health supervisor

d) village health guide
169) Least recurrence of peptic ulcer most commonly occurs in
a) selective vagotomy
b) TV + GJ
c) Pure gastroenterostomy
d) TV+ antrectomy
170) CABG has successful outcome and prognosis if done in
a) double vessel obstruction
b) left coronary artery obstruction without left ventricular dysfunction
c) single vessel obstruction
171) anteversion of uterus is due to following ligament
a) uterosacral ligament
b) mackenrod's ligament
c)
d)
172) osteochondroma is a disease of
a)metaphysis
b) diaphysis
c) epiphysis

d) periosteum
173) the characteristic symptom of dysmenorrhea , menorrhagia, recurrent abdominal pain and infertility is suggestive of
a) endometriosis
b) ectopic
c) endometritis
d)
174) testicular seminoma, secretes
a) ? -FP
b) CEA
c) B-HCG
d)
175) the concentration of HbA1c in blood is
a) 4-6%
b) 6-8%
c) 8-9%
d)
176) which of the following is contraindicated in early pregnancy
a) heparin
b) phenytoin

c) chlor***
d) penicillin
177) action potential is transmitted to myofibrils via
a) terminal cisterns
b) T- Tubules
c) longitudinal
d) sarcomere
178) humidification of O2 is required in
a) nasopharyngeal intubation
b) nasal prong
c) nasal catheter
d) face mask
179) meconium plug syndrome occurs in
a) meconium ileus
b) hirschsprung's disease
c) ileo-colic pouch
d)
180) the cause of endemic ascites is
a) BOAA

b) Sanguinarine
c) Pyrozzilodine alkaloids
d)
181) earliest sign of vitamin A deficiency is
a) conjunctival xerosis
b) corneal xerosis
c) bitot's spot
d)nightblindness
182) Pinta is caused by
a) T pertenue
b) T pallidum
c) T carateum
d) Borrelia burgdorferi.
183) all the following cause priapism except
a) trauma
b) Beta thalassemia
c) sickle cell disease
d) leriche syndrome
184) iridectomy is done in which type of glaucoma is

a)
c) angle closure
b)
d) open glaucoma
185) yellow fever is not present in India because of all except
a) the virus is absent
b) the vector mosquito culex vishnui
c) atmospheric condition is suitable
186) phage typing is useful in classification of all except
a) staphylococcus
b) streptococcus
c) cholera
d) salmonella
187)in a patient with strikingly abnormal behavior, the most likely diagnosis is
a) paranoid schizophrenia
b) catatonic schizophrenia
c) hebephrenic schizophrenia
d)
188)the revised national TB program uses one of the following for diagnosis of TB

a) sputum AFB
b) CXR
c) clinical examination
d) PCR
189) following are true of rubella except
a) IP: 14-21days
b) rosy red spots on palate
c) encephalitis
190) Acrodynia occurs in
a) mercury poisoning
b) arsenic poisoning
c) lead poisoning
d)
191) hydrotherapy is done in
a) impotence
b) infertility
c) febrile convulsion
d)

192) hemodialysis is useful in all except

a) salicylate poisoning
b) methyl alcohol poisoning
c) digitalis
d) barbiturate
193) proliferation of mesangial cell & perimesangial on microscopy is seen in
a) lg A nephropathy
b) lipoid nephrosis
c) membranoproliferative disease
d) membranous glomerulonephritis
194) point prevalence of a chronic disease is studied best by
a) case control study
b) cross sectional study
c) historical cohort
d) cohort
195) reversible cataract is seen in which of following
a) galactosemia
b) hypoparathyroidism
c) diabetes mellitus
d)

196) De quervain's thyroiditis is due to all except
a) it is due to virus
b) antibodies against follicular cells
c) it is self limited granulomatous thyroiditis
197)partial prothromboplastin time is prolonged in all except
a) protein C deficiency
b) protein S deficiency
c) antithrombin III deficiency
d) factor V deficiency
198) which of the following has least t1/2
a) norepinephrine
b) adrenaline
c) renin
d)
199) the treatment of primary ovarian failure is
a) FSH
b) GnRH
c)LH
d)estrogen progesterone
200)LDL cholesterol in blood recommended to prevent coronary artery disease is

- a)130mg/dl
- b)160mg/dl
- c)100mg/dl
- d)180mg/dl